

Julie Estes-Undergraduate Researcher¹, Dr. Bart Tarbet-Faculty Advisor²

¹Department of Biology – College of Science – Utah State University

²Department of Animal Dairy and Veterinarian Sciences



INTRODUCTION

Our goal is to develop diagnostic assays for the viruses associated with Bovine Respiratory Disease (BRD). BRD is a respiratory disease and is the result of five different viruses including Bovine Viral Diarrhea Virus (BVDV) types 1 and 2, infectious bovine rhinotracheitis, bovine parainfluenza-3 viruses, and bovine respiratory syncytial virus. BVDV causes immunosuppression of the infected adaptive immune system. Our main objective is to reduce the loss of the cattle population due to this virus. The ability to control and eradicate BVDV would reduce the significant economic loss of the cattle population. [1] Our project is to develop diagnostic procedures that will detect the infection in its many forms.



Figure 2 (right) - Julie Estes in lab

.METHODS.

I have been learning certain scientific techniques that will be imperative in continuing this research. They include:

- Cell culturing
The process of growing cells under controlled conditions.
- Cell Seeding 96 Well Plate
A standard tool in analytical research and clinical diagnostic testing.
- Hemocytometer
Allowance of selective cell counting
- Virology
Replication of viruses in cell culture for use in clinical detection and isolation of viruses, in addition to how they grow and infect organisms.



Figure 3 - Stock image of seeding 96 well plate

BVDV is currently one of the most costly diseases of cattle. Abortions, infertility, and/or embryonic deaths associated with...

...BVDV lead to significantly reduced reproductive performance and increased premature culling. [2]

.RESULTS.

(from earlier studies)

“Using viral neutralization assays, we found vital evaluation parameters were cell line, virus concentration, and incubation time. Using the establish protocol above we found that the average of the numerical protection average from the serums tested was around 169.24. An average standard deviation of 64.65 was obtained, which is a 33 % of test varied by about one dilution step from trial to trial.”
-Adam Allen, Utah State University

FUTURE STUDIES

- Evaluate the effect of different bovine serum dilutions
- Qualifying viral neutralization assay protocols for use with BVDV.
- Testing cell lines, virus concentrations, and incubation variability for assay

ACKNOWLEDGEMENTS

Special thanks to faculty and personnel from the Institute for Antiviral Research, specifically Rebecca Strong for the training and collaboration.

REFERENCES

- [1] Dubovi, E. (2012). Laboratory diagnosis of bovine viral diarrhoea virus, www.elsevier.com/locate/biologicals
[2] Animal Health Diagnostic Center, Cornell University (2016). Bovine Viral Diarrhoea: Background, Management and Control, <https://www.vet.cornell.edu/>